

## Flu eligibility criteria

The eligibility criteria for the Autumn 2025 flu vaccination campaign is:

- those aged 65 years and over
- those in long-stay residential care homes
- those aged 18 years to under 65 years in clinical risk groups as defined by the table below
- carers in receipt of carer's allowance, or those who are the main carer of an elderly or disabled person
- close contacts of immunocompromised individuals
- frontline workers in a social care setting without an employer led occupational health scheme including those working for a registered residential care or nursing home, registered domiciliary care providers, voluntary managed hospice providers and those that are employed by those who receive direct payments (personal budgets) or Personal Health budgets, such as Personal Assistants

Clinical risk category	Examples (this list is not exhaustive and decisions should be based on clinical judgement)
Chronic respiratory disease	Asthma that requires continuous or repeated use of inhaled or systemic steroids or with previous exacerbations requiring hospital admission. Chronic obstructive pulmonary disease (COPD) including chronic bronchitis and emphysema; bronchiectasis, cystic fibrosis, interstitial lung fibrosis, pneumoconiosis and bronchopulmonary dysplasia (BPD). In addition to those with chronic respiratory disease, children who have previously been admitted to hospital for lower respiratory tract disease. See precautions section on LAIV.
Chronic heart disease and vascular disease	Congenital heart disease, hypertension with cardiac complications, chronic heart failure, individuals requiring regular medication and/or follow-up for ischaemic heart disease. This includes individuals with atrial fibrillation, peripheral vascular disease or a history of venous thromboembolism.
Chronic kidney disease	Chronic kidney disease at stage 3, 4 or 5, chronic kidney failure, nephrotic syndrome, kidney transplantation.
Chronic liver disease	Cirrhosis, biliary atresia, chronic hepatitis.
Chronic neurological disease (included in the DES directions for Wales)	Stroke, transient ischaemic attack (TIA). Conditions in which respiratory function may be compromised due to neurological or neuromuscular disease (for example polio syndrome sufferers). Clinicians should offer immunisation, based on individual assessment, to clinically vulnerable individuals including those with cerebral palsy, severe or profound and multiple learning disabilities (PMLD), Down's syndrome, multiple sclerosis, dementia, Parkinson's disease, motor neurone disease and related or similar conditions; or hereditary and degenerative disease of the nervous system or muscles; or severe neurological disability.
Diabetes and adrenal insufficiency	Type 1 diabetes, type 2 diabetes requiring insulin or oral hypoglycaemic drugs, diet-controlled diabetes. Addison's disease, secondary or tertiary adrenal insufficiency requiring steroid replacement.

<b>Clinical risk category</b>	<b>Examples (this list is not exhaustive and decisions should be based on clinical judgement)</b>
Immunosuppression (see contraindications and precautions section on live attenuated influenza vaccine)	<p>Immunosuppression due to disease or treatment, including patients undergoing chemotherapy leading to immunosuppression, patients undergoing radical radiotherapy, solid organ transplant recipients, bone marrow or stem cell transplant recipients, people living with HIV (at all stages), multiple myeloma or genetic disorders affecting the immune system (for example IRAK-4, NEMO, complement disorder, SCID). Individuals who are receiving immunosuppressive or immunomodulating biological therapy including, but not limited to, anti-TNF- alemtuzumab, ofatumumab, rituximab, patients receiving protein kinase inhibitors or PARP inhibitors, and individuals treated with steroid sparing agents such as cyclophosphamide and mycophenolate mofetil.</p> <p>Individuals treated with or likely to be treated with systemic steroids for more than a month at a dose equivalent to prednisolone at 20mg or more per day (any age), or for children under 20kg, a dose of 1mg or more per kg per day.</p> <p>Anyone with a history of haematological malignancy, including leukaemia, lymphoma, and myeloma and those with systemic lupus erythematosus and rheumatoid arthritis, and psoriasis who may require long term immunosuppressive treatments.</p> <p>Some immunocompromised patients may have a suboptimal immunological response to the vaccine.</p>
Asplenia or dysfunction of the spleen	This also includes conditions such as homozygous sickle cell disease, hereditary spherocytosis, thalassemia major and coeliac disease that may lead to splenic dysfunction.
Morbid obesity (class III obesity)*	Adults with a Body Mass Index $\geq 40$ kg/m <sup>2</sup> .

<b>Other risk groups</b>	
Pregnant women	Pregnant women at any stage of pregnancy (first, second or third trimesters). See precautions section on live attenuated influenza vaccine.
Household contacts of people with immunosuppression	Individuals who expect to share living accommodation on most days (and therefore for whom continuing close contact is unavoidable) with individuals who are immunosuppressed (defined as immunosuppressed in table 19.4).
Carers	Those who are eligible for a carer's allowance, or those who are the sole or primary carer of an elderly or disabled person whose welfare may be at risk if the carer falls ill.

\* Many of this patient group will already be eligible due to complications of obesity that place them in another risk category

The list above is not exhaustive, and the medical practitioner should apply clinical judgment to take into account the risk of influenza exacerbating any underlying disease that a patient may have, as well as the risk of serious illness from influenza itself. Influenza vaccine should be offered in such cases even if the individual is not in the clinical risk groups specified above.