

Flu vaccination eligibility 2026/2027

Influenza vaccine should be offered to:

- all those aged 65 years or older
- all those aged 6 months or older in the clinical risk groups shown in tables below
- all children in cohorts eligible for vaccination as part of the children's flu programme (see below)
- those living in long-stay residential care homes or other long-stay care facilities where rapid spread is likely to follow introduction of infection and cause high morbidity and mortality (this does not include prisons, young offender institutions, university halls of residence etc.)

Table 19.4 Clinical risk groups and other risk groups who should be offered influenza vaccination.

Clinical risk category	Examples (this list is not exhaustive and decisions should be based on clinical judgement)
Chronic respiratory disease	Asthma that requires continuous or repeated use of inhaled or systemic steroids or with previous exacerbations requiring hospital admission. Chronic obstructive pulmonary disease (COPD) including chronic bronchitis and emphysema; bronchiectasis, cystic fibrosis, interstitial lung fibrosis, pneumoconiosis and bronchopulmonary dysplasia (BPD). In addition to those with chronic respiratory disease, children who have previously been admitted to hospital for lower respiratory tract disease. See precautions section on LAIV.
Chronic heart disease and vascular disease	Congenital heart disease, hypertension with cardiac complications, chronic heart failure, individuals requiring regular medication and/or follow-up for ischaemic heart disease. This includes individuals with atrial fibrillation, peripheral vascular disease or a history of venous thromboembolism.
Chronic kidney disease	Chronic kidney disease at stage 3, 4 or 5, chronic kidney failure, nephrotic syndrome, kidney transplantation.
Chronic liver disease	Cirrhosis, biliary atresia, chronic hepatitis.
Chronic neurological disease (included in the DES directions for Wales)	Stroke, transient ischaemic attack (TIA). Conditions in which respiratory function may be compromised due to neurological or neuromuscular disease (for example polio syndrome sufferers). Clinicians should offer immunisation, based on individual assessment, to clinically vulnerable individuals including those with cerebral palsy, severe or profound and multiple learning disabilities (PMLD), Down's syndrome, multiple sclerosis, dementia, Parkinson's disease, motor neurone disease and related or similar conditions; or hereditary and degenerative disease of the nervous system or muscles; or severe neurological disability.
Diabetes and adrenal insufficiency	Type 1 diabetes, type 2 diabetes requiring insulin or oral hypoglycaemic drugs, diet-controlled diabetes. Addison's disease, secondary or tertiary adrenal insufficiency requiring steroid replacement.

Clinical risk category	Examples (this list is not exhaustive and decisions should be based on clinical judgement)
Immunosuppression (see contraindications and precautions section on live attenuated influenza vaccine)	Immunosuppression due to disease or treatment, including patients undergoing chemotherapy leading to immunosuppression, patients undergoing radical radiotherapy, solid organ transplant recipients, bone marrow or stem cell transplant recipients, people living with HIV (at all stages), multiple myeloma or genetic disorders affecting the immune system (for example IRAK-4, NEMO, complement disorder, SCID). Individuals who are receiving immunosuppressive or immunomodulating biological therapy including, but not limited to, anti-TNF-α (adalimumab, infliximab, certolizumab, etanercept), rituximab, patients receiving protein kinase inhibitors or PARP inhibitors, and individuals treated with steroid sparing agents such as cyclophosphamide and mycophenolate mofetil. Individuals treated with or likely to be treated with systemic steroids for more than a month at a dose equivalent to prednisolone at 20mg or more per day (any age), or for children under 20kg, a dose of 1mg or more per kg per day. Anyone with a history of haematological malignancy, including leukaemia, lymphoma, and myeloma and those with systemic lupus erythematosus and rheumatoid arthritis, and psoriasis who may require long term immunosuppressive treatments. Some immunocompromised patients may have a suboptimal immunological response to the vaccine.
Asplenia or dysfunction of the spleen	This also includes conditions such as homozygous sickle cell disease, hereditary spherocytosis, thalassemia major and coeliac disease that may lead to splenic dysfunction.
Morbid obesity (class III obesity)*	Adults with a Body Mass Index ≥ 40 kg/m ² .

Other risk groups	
Pregnant women	Pregnant women at any stage of pregnancy (first, second or third trimesters). See precautions section on live attenuated influenza vaccine.
Household contacts of people with immunosuppression	Individuals who expect to share living accommodation on most days (and therefore for whom continuing close contact is unavoidable) with individuals who are immunosuppressed (defined as immunosuppressed in table 19.4).
Carers	Those who are eligible for a carer's allowance, or those who are the sole or primary carer of an elderly or disabled person whose welfare may be at risk if the carer falls ill.
People experiencing homelessness	Rough sleepers and those using homeless hostels or night-shelters

* Many of this patient group will already be eligible due to complications of obesity that place them in another risk category